

New class of multiple weights and new weighted inequalities for multilinear operators

The Anh Bui*

Abstract

In this paper, we first introduce the new class of multiple weights $A_{\vec{p}}^{\infty}$ which is larger than the class of multiple weights in [LOPTG]. Then, using this class of weights, we study the weighted norm inequalities for certain classes of multilinear operators and their commutators with new BMO functions introduced by [BHS1]. Finally, we show that some multilinear pseudodifferential operators fall within the scope of the theory obtained in this paper.

Contents

1	Introduction and the main results	1
2	Preliminaries	3
2.1	The new class of weights and new BMO function spaces	3
2.1.1	Classes of multiple weights $A_{\vec{p}}^{\infty}$	3
2.1.2	New BMO function spaces BMO_{∞}	6
2.2	Weighted estimates for some maximal operators	6
3	Main results	10
3.1	Weighted norm inequalities for multilinear operators	10
3.2	Weighted norm inequalities for the commutators of multilinear operators with BMO_{∞} functions	13
4	Application to multilinear pseudodifferential operators	17

1 Introduction and the main results

The theory of multilinear Calderón-Zygmund singular integral operators, originated from the work of Coifman and Meyer, has an important role in harmonic analysis. This direction of research has been attracting a lot of attention in the last few decades, see for example [CM1, CM2, CM3, GT, KS] for the standard theory of multilinear Calderón-Zygmund singular integrals.

*Department of Mathematics, Macquarie University, NSW 2109, Australia and Department of Mathematics, University of Pedagogy, HoChiMinh City, Vietnam.

Email: the.bui@mq.edu.au and bt_anh80@yahoo.com

2010 *Mathematics Subject Classification*: 42B20, 42B25, 35S05, 47G30.

Key words: Weighted norm inequality; Multilinear operator; Multilinear pseudodifferential; Commutator.

Let T be a multilinear operator initially defined on the m -fold product of Schwartz spaces and taking values into the space of tempered distributions,

$$T : \mathcal{S}(\mathbb{R}^n) \times \dots \times \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n).$$

By associated kernel to T we shall mean the function K , defined off the diagonal $x = y_1 = \dots = y_m$ in $(\mathbb{R}^n)^{m+1}$, satisfying

$$T(f_1, \dots, f_m)(x) = \int_{(\mathbb{R}^n)^m} K(x, y_1, \dots, y_m) f_1(y_1) \dots f_m(y_m) dy_1 \dots dy_m$$

for all $x \notin \cap_{j=1}^m \text{supp } f_j$.

In this paper, we consider the following conditions:

(H1) For any $N > 0$ there exists $C > 0$ such that

$$|K(y_0, y_1, \dots, y_m)| \leq \frac{C}{(\sum_{k,l=0}^m |y_k - y_l|)^N}. \quad (1)$$

(H2) For any $N > 0$ there exists $C > 0$ such that

$$|K(y_0, \dots, y_j, \dots, y_m) - K(y_0, \dots, y'_j, \dots, y_m)| \leq \frac{C|y_j - y'_j|^\epsilon}{(\sum_{k,l=0}^m |y_k - y_l|)^{mn+\epsilon}} \min\{1, h^{-N}\}, \quad (2)$$

for some $\epsilon > 0$ and all $0 \leq j \leq m$, whenever $|y_j - y'_j| \leq \frac{1}{2} \max_{0 \leq k \leq m} |y_j - y_k| := h$.

(H3) There exist $1 \leq q_1, \dots, q_m < \infty$ and $1/q = 1/q_1 + \dots + 1/q_m$ such that T maps continuously from $L^{q_1} \times \dots \times L^{q_m}$ into L^q .

It is clear that if T satisfies (H1), (H2) and (H3) then T falls within the scope of multilinear Calderón-Zygmund theory investigated by [GT]. Therefore, according to [GT], if $1/p = 1/p_1 + \dots + 1/p_m$, the following statements hold:

- (i) $T : L^{p_1} \times \dots \times L^{p_m} \rightarrow L^p$ when $1 < p_1, \dots, p_m < \infty$, and
- (ii) $T : L^{p_1} \times \dots \times L^{p_m} \rightarrow L^{p,\infty}$ when $1 \leq p_1, \dots, p_m < \infty$ and at least one $p_j = 1$.

The weighted norm inequalities of multilinear Calderón-Zygmund operators and their commutators with BMO functions were investigated in [LOPTG]. In [LOPTG], the authors introduced the new maximal functions and multiple weights and then they proved that the new class of multiple weights is suitable to study the boundedness of multilinear Calderón-Zygmund operators and their commutators with BMO functions.

Inspiring by the works of [LOPTG] for boundedness of multilinear Calderón-Zygmund operators and their commutators with BMO functions and of [BHS1, BHS2] for the new class of weights and new BMO function spaces, the aim of this paper is to study the weighted norm inequalities of operators T which satisfy (H1)-(H3) and their commutators by using the new BMO function spaces introduced by [BHS1] and the new class of multiple weights introduced in Section 2.

The organization of the paper is as follows. In Section 2, we introduce the new class of multiple weights and then investigate the weighted norm inequalities of some maximal functions. Section 3 establishes the main results of the paper. Firstly, the weighted estimates of multilinear

operators T are investigated (see Theorem 3.1). Secondly, we consider the weighted norm inequalities of the commutator T_b by using the new BMO functions and new class of multiple weights (see Theorem 3.6). In Section 4, we show that the obtained results can be applied to certain multilinear pseudodifferential operators.

After finishing this paper, I had informed that the author in [T] obtained the A_p^∞ weighted norm inequalities for such an operator T . However, in this paper, we study the the A_p^∞ weighted norm inequalities of T and the obtained results and the new class of multiple weights in our paper are new. Moreover, the weighted norm inequalities of the commutator T_b with the new BMO functions in our paper are unique.

2 Preliminaries

To simplify notation, we will often just use B for $B(x_B, r_B)$ and $|E|$ for the measure of E for any measurable subset $E \subset \mathbb{R}^n$. Also given $\lambda > 0$, we will write λB for the λ -dilated ball, which is the ball with the same center as B and with radius $r_{\lambda B} = \lambda r_B$. For each ball $B \subset \mathbb{R}^n$ we set

$$S_0(B) = B \text{ and } S_j(B) = 2^j B \setminus 2^{j-1} B \text{ for } j \in \mathbb{N}.$$

2.1 The new class of weights and new BMO function spaces

2.1.1 Classes of multiple weights A_p^∞

In this section, we would like to recall the definition of the new class of weights introduced by [BHS2].

For $1 \leq p < \infty$ and $\theta \geq 0$, the weight w (w is a nonnegative and locally integrable function) is said to be in the class A_p^θ if there holds

$$\left(\int_B w \right)^{1/p} \left(\int_B w^{-\frac{1}{p-1}} \right)^{1/p'} \leq C|B|(1+r_B)^\theta \quad (3)$$

for all ball $B = B(x_B, r_B)$. In particular case when $p = 1$, (3) is understood

$$\frac{1}{|B|} \int_B w(y) dy \leq C(1+r_B)^\theta \inf_{x \in B} w(x).$$

Then we denote $A_p^\infty = \cup_{\theta \geq 0} A_p^\theta$ and $A_\infty^\infty = \cup_{p \geq 1} A_p^\infty$.

We remark that A_p^0 coincides with the Muckenhoupt's class of weights A_p for all $1 \leq p < \infty$. However, in general, the class A_p^∞ is strictly larger than the class A_p for all $1 \leq p < \infty$. The following properties hold for the new classes A_p^∞ , see [BHS2, Proposition 5].

Proposition 2.1 *The following statements hold:*

- i) $A_p^\infty \subset A_q^\infty$ for $1 \leq p \leq q < \infty$.
- ii) If $w \in A_p^\infty$ with $p > 1$ then there exists $\epsilon > 0$ such that $w \in A_{p-\epsilon}^\infty$. Consequently, $A_p^\infty = \cup_{q < p} A_q^\infty$.
- iii) If $w \in A_p^\infty$ with $p \geq 1$, then there exist positive numbers δ, η and C so that for all balls B ,

$$\left(\frac{1}{|B|} \int_B w^{1+\delta}(x) dx \right)^{\frac{1}{1+\delta}} \leq C \left(\frac{1}{|B|} \int_B w(x) dx \right) (1+r_B)^\eta.$$

In [LOPTG], to study the weighted norm inequalities of multilinear operators, the authors introduced the new maximal functions and the multiple weights. Adapting this idea to our situation, we introduce the new class of multiple weights.

In what follows, for given m exponents p_1, \dots, p_m , unless specified, otherwise we write $\vec{p} = (p_1, \dots, p_m)$ and the number p shall mean that

$$\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}.$$

For any number $r > 0$, $r\vec{p}$ is defined by $r\vec{p} = (rp_1, \dots, rp_m)$.

Definition 2.2 Let $1 \leq p_1, \dots, p_m < \infty$. For $\vec{w} = (w_1, \dots, w_m)$, set

$$\nu_{\vec{w}} = \prod_{j=1}^m w_j^{p/p_j}.$$

For $\theta \geq 0$, we say that \vec{w} is in the class $A_{\vec{p}}^{\theta}$ if

$$\left(\frac{1}{|B|} \int_B \nu_{\vec{w}}(x) dx \right)^{1/p} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{1-p'_j}(x) dx \right)^{1/p'_j} \leq C(1+r_B)^{\theta}$$

for all balls B . When $p_j = 1$, $\left(\frac{1}{|B|} \int_B w_j^{1-p'_j}(x) dx \right)^{1/p'_j}$ is understood $(\inf_{x \in Q} w_j(x))^{-1}$.

For $1 \leq p_1, \dots, p_m < \infty$, we set $A_{\vec{p}}^{\infty} = \cup_{\theta \geq 0} A_{\vec{p}}^{\theta}$.

When $\theta = 0$, the class $A_{\vec{p}}^0$ coincides with the class of multiple weights $A_{\vec{p}}$ introduced by [LOPTG]. The following result gives a characterization of the class $A_{\vec{p}}^{\infty}$ whose proof is similar to that of [LOPTG, Theorem 3.6].

Proposition 2.3 Let $1 \leq p_1, \dots, p_m < \infty$ and $\vec{w} = (w_1, \dots, w_m)$. Then the following statements are equivalent:

- (i) $\vec{w} \in A_{\vec{p}}^{\infty}$;
- (ii) $w_j^{1-p'_j} \in A_{mp'_j}^{\infty}, j = 1, \dots, m$ and $\nu_{\vec{w}} \in A_{mp}^{\infty}$.

Note that the class $A_{\vec{p}}^{\infty}$ is not increasing. It means that for $\vec{p} = (p_1, \dots, p_m)$ and $\vec{q} = (q_1, \dots, q_m)$ with $p_j \leq q_j, j = 1, \dots, m$, the following may not be true $A_{\vec{p}}^{\infty} \subset A_{\vec{q}}^{\infty}$, see [LOPTG, Remark 7.3]. However, we have the following result.

Proposition 2.4 Let $1 \leq p_1, \dots, p_m < \infty$ and $\vec{w} = (w_1, \dots, w_m) \in A_{\vec{p}}^{\infty}$. Then,

- (i) For any $r \geq 1$, $\vec{w} \in A_{r\vec{p}}^{\infty}$;
- (ii) If $1 < p_1, \dots, p_m < \infty$, then there exists $r > 1$ so that $\vec{w} \in A_{\vec{p}/r}^{\infty}$.

Proof: (i) Assume that $\vec{w} \in A_{\vec{p}}^{\theta}$ for some $\theta \geq 0$. By definition, there exists $C > 0$ so that for all balls B there holds,

$$\left(\frac{1}{|B|} \int_B \nu_{\vec{w}}(x) dx \right)^{1/p} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{p_j-1}}(x) dx \right)^{\frac{p_j-1}{p_j}} \leq C(1+r_B)^{\theta}.$$

For $r > 1$, by Hölder inequality we have

$$\left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{rp_j-1}}(x) dx \right)^{\frac{rp_j-1}{rp_j}} \leq \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{p_j-1}}(x) dx \right)^{\frac{p_j-1}{rp_j}}.$$

Therefore, for all balls B , we have

$$\left(\frac{1}{|B|} \int_B \nu_{\vec{w}}(x) dx \right)^{1/rp} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{rp_j-1}}(x) dx \right)^{\frac{rp_j-1}{rp_j}} \leq C(1+r_B)^{\theta/r}.$$

This implies that $\vec{w} \in A_{r\vec{p}}^\infty$.

(ii) We exploit some ideas in [LOPTG] to our situation. In the light of Propositions 2.1 and 2.3, we can pick $r_0 > 1$ and $\eta \geq 0$ so that

$$\left(\frac{1}{|B|} \int_B w_j^{-\frac{r_0}{p_j-1}}(x) \right)^{1/r_0} \leq C \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{p_j-1}}(x) \right) (1+r_B)^\theta \quad (4)$$

for all balls B and $j = 1, \dots, m$.

Taking $r > 1$ so that $r < \frac{r_0 p_j}{p_j + r_0 - 1}$ for all $j = 1, \dots, m$, then we have for all j ,

$$\frac{r_0(p_j - r)}{r(p_j - 1)} < 1.$$

This together with Hölder inequality gives

$$\left(\frac{1}{|B|} \int_B w_j^{-\frac{r}{p_j-r}}(x) dx \right)^{\frac{p_j-r}{p_j}} \leq \left(\frac{1}{|B|} \int_B w_j^{-\frac{r_0}{p_j-1}}(x) dx \right)^{\frac{r(p_j-1)}{r_0 p_j}}.$$

Due to (4), we have

$$\left(\frac{1}{|B|} \int_B w_j^{-\frac{r}{p_j-r}}(x) dx \right)^{\frac{p_j-r}{p_j}} \leq \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{p_j-1}}(x) dx \right)^{\frac{r(p_j-1)}{p_j}} (1+r_B)^{r\theta}.$$

Therefore,

$$\left(\frac{1}{|B|} \int_B \nu_{\vec{w}}(x) dx \right)^{r/p} \prod_{j=1}^m \left(\frac{1}{|B|} \int_B w_j^{-\frac{1}{p_j/r-1}}(x) dx \right)^{\frac{p_j/r-1}{p_j/r}} \leq C(1+r_B)^{r\theta}.$$

It yields $\vec{w} \in A_{\vec{p}/r}^\infty$.

□

2.1.2 New BMO function spaces BMO_∞

In this section, we will recall the definition and some basic properties of the new BMO function spaces. According to [BHS1], the new BMO space BMO_θ with $\theta \geq 0$ is defined as a set of all locally integrable functions b satisfying

$$\frac{1}{|B|} \int_B |b(y) - b_B| dy \leq C(1 + r_B)^\theta \quad (5)$$

where $B = B(x_B, r_B)$ and $b_B = \frac{1}{|B|} \int_B b$. A norm for $b \in BMO_\theta$, denoted by $\|b\|_\theta$, is given by the infimum of the constants satisfying (5). Clearly $BMO_{\theta_1} \subset BMO_{\theta_2}$ for $\theta_1 \leq \theta_2$ and $BMO_0 = BMO$. We define $BMO_\infty = \cup_{\theta \geq 0} BMO_\theta$.

The following result can be considered to be a variant of John-Nirenberg inequality for the spaces BMO_∞ .

Proposition 2.5 *Let $\theta > 0, s \geq 1$. If $b \in BMO_\theta$ then for all $B = (x_0, r)$*

i)

$$\left(\frac{1}{|B|} \int_B |b(y) - b_B|^s dx \right)^{1/s} \lesssim \|b\|_\theta (1 + r_B)^\theta;$$

ii)

$$\left(\frac{1}{|2^k B|} \int_{2^k B} |b(y) - b_B| dx \right)^{1/s} \lesssim \|b\|_\theta k (1 + 2^k r_B)^\theta$$

for all $k \in \mathbb{N}$.

We refer to Proposition 3 in [BHS2] for the proof.

2.2 Weighted estimates for some maximal operators

A ball of the form $B(x_B, r_B)$ is called a *critical ball* if $r_B = 1$. We have the following result.

Proposition 2.6 *There exists a sequence of points $x_j, j \geq 1$ in \mathbb{R}^n so that the family of critical balls $\{Q_j\}_j$ where $Q_j := B(x_j, 1)$, $j \geq 1$ satisfies*

(i) $\cup_j Q_j = \mathbb{R}^n$.

(ii) There exists a constant C such that for any $\sigma > 1$, $\sum_j \chi_{\sigma Q_j} \leq C\sigma^n$.

For the proof, we refer the reader to [B] (see also [DZ]).

We consider the following maximal functions for $g \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$

$$M_{\text{loc}, \alpha} g(x) = \sup_{x \in B \in \mathcal{B}_\alpha} \frac{1}{|B|} \int_B |g|,$$

$$M_{\text{loc}, \alpha}^\# g(x) = \sup_{x \in B \in \mathcal{B}_\alpha} \frac{1}{|B|} \int_B |g - g_B| \approx \inf_{c \in \mathbb{R}} \sup_{x \in B \in \mathcal{B}_\alpha} \frac{1}{|B|} \int_B |g - c|,$$

where $\mathcal{B}_\alpha = \{B(y, r) : y \in \mathbb{R}^n \text{ and } r \leq \alpha\}$.

Also, given a ball Q , we define the following maximal functions for $g \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $x \in Q$

$$M_Q g(x) = \sup_{x \in B \in \mathcal{F}(Q)} \frac{1}{|B \cap Q|} \int_{B \cap Q} |g|,$$

$$M_Q^\sharp g(x) = \sup_{x \in B \in \mathcal{F}(Q)} \frac{1}{|B \cap Q|} \int_{B \cap Q} |g - g_{B \cap Q}|,$$

where $\mathcal{F}(Q) = \{B(y, r) : y \in Q, r > 0\}$.

We have the following lemma.

Lemma 2.7 *For $0 < p < \infty$, then there exists β such that if $\{Q_k\}_k$ is a sequence of balls as in Proposition 2.6 then for all $g \in L_{\text{loc}}^1(\mathbb{R}^n)$ and $w \in A_\infty^\infty$, we have*

(i)

$$\int_{\mathbb{R}^n} |M_{\text{loc}, \beta} g(x)|^p w(x) dx \lesssim \int_{\mathbb{R}^n} |M_{\text{loc}, 4}^\sharp g(x)|^p w(x) dx + \sum_k w(2Q_k) \left(\frac{1}{|2Q_k|} \int_{2Q_k} |g| \right)^p;$$

and

$$(ii) \|M_{\text{loc}, \beta} g\|_{L^{p, \infty}(w)}^p \leq \|M_{\text{loc}, 4}^\sharp g\|_{L^{p, \infty}(w)}^p + \sum_k w(2Q_k) \left(\frac{1}{|2Q_k|} \int_{2Q_k} |g| \right)^p.$$

Proof: We refer to [B, Lemma 2.4] for the proof of (i). The proof of (ii) is similar to that of (i) and we omit details here. □

Throughout this paper, we always assume that N is a sufficiently large number and different from line to line. For $\kappa \geq 1$, $p > 0$, $\vec{f} = (f_1, \dots, f_m)$, $f_j \in L_{\text{loc}}^1(\mathbb{R}^n)$ for all $j = 1, \dots, m$ and $x \in \mathbb{R}^n$, we define the maximal function \mathcal{M} by

$$\mathfrak{M}_{\kappa, p}(\vec{f})(x) = \sup_{Q \ni x; Q \text{ is critical}} \sum_{k=0}^{\infty} 2^{-Nk} \prod_{j=1}^m \left(\frac{1}{|2^k \widehat{Q}|} \int_{2^k \widehat{Q}} |f_j(z)|^p dz \right)^{1/p}$$

where $\widehat{Q} = \kappa Q$.

For simplicity, we shall write \mathfrak{M}_p and \mathfrak{M} instead of $\mathfrak{M}_{1, p}$ and $\mathfrak{M}_{1, 1}$, respectively. The following result gives the weighted estimates for $\mathfrak{M}_{\kappa, p}$.

Proposition 2.8 *Let $p_1, \dots, p_m \geq s > 0$ and $w \in A_{\vec{p}/s}^\infty$. Then we have*

$$\|\mathfrak{M}_{\kappa, s}(\vec{f})\|_{L^p(\nu_{\vec{w}})} \lesssim \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

Proof: Without of the loss of generality, we assume that $\alpha = 1$. Let $\{Q_\ell\}$ be the family of balls as in Proposition 2.6. Then we have

$$\|\mathfrak{M}_s(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \left(\sum_\ell \int_{Q_\ell} |\mathfrak{M}_s(\vec{f})(x)|^p \nu_{\vec{w}} dx \right)^{1/p}.$$

If $x \in Q_\ell \cap Q$, where Q is a critical ball, then $2^k Q \subset 2^{k+1} Q_\ell$ and $|2^k Q| \approx |2^{k+1} Q_\ell|$. Therefore, for $x \in Q_\ell$ we have

$$\mathfrak{M}_s(\vec{f})(x) \leq C \sum_{k=0}^{\infty} 2^{-Nk} \prod_{j=1}^m \left(\frac{1}{|2^k Q_\ell|} \int_{2^k Q_\ell} |f_j(z)|^s dz \right)^{1/s}.$$

So,

$$\begin{aligned}
\|\mathfrak{M}_s(\vec{f})\|_{L^p(\nu_{\vec{w}})} &\leq C \left(\sum_{\ell} \int_{Q_{\ell}} \left| \sum_{k=0}^{\infty} 2^{-Nk} \prod_{j=1}^m \left(\frac{1}{|2^k Q_{\ell}|} \int_{2^k Q_{\ell}} |f_j(z)|^s dz \right)^{1/s} \right|^p \nu_{\vec{w}} dx \right)^{1/p} \\
&\leq C \sum_{k=0}^{\infty} 2^{-Nk} \left(\sum_{\ell} \nu_{\vec{w}}(Q_{\ell}) \prod_{j=1}^m \left(\frac{1}{|2^k Q_j|} \int_{2^k Q_j} |f_j(z)|^s dz \right)^{p/s} \right)^{1/p} \\
&\leq C \sum_{k=0}^{\infty} 2^{-Nk} \left(\sum_{\ell} \nu_{\vec{w}}(2^k Q_{\ell}) \prod_{j=1}^m \left(\frac{1}{|2^k Q_j|} \int_{2^k Q_j} |f_j(z)|^s dz \right)^{p/s} \right)^{1/p}.
\end{aligned}$$

Assume that $\vec{w} \in A_{\vec{p}/s}^{\theta}$ for some $\theta \geq 0$. For each j , by Hölder inequality, we have

$$\int_{2^k Q_{\ell}} |f_j(z)|^s dz \leq \|f_j\|_{L^{p_j}(w_j, 2^k Q_{\ell})}^s \left(\int_{2^k Q_{\ell}} w^{-\frac{1}{p_j/s-1}} \right)^{\frac{p_j/s}{p_j/s-1}}.$$

This together with definition of $A_{\vec{p}}^{\theta}$ class gives

$$\|\mathfrak{M}_s(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \sum_{k=0}^{\infty} 2^{-k(N-\theta/s)} \left(\sum_{\ell} \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j, 2^k Q_{\ell})}^p \right)^{1/p}.$$

Using Hölder inequality and (ii) of Proposition 2.6, we get that

$$\begin{aligned}
\sum_{\ell} \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j, 2^k Q_{\ell})}^p &\leq \prod_{j=1}^m \left(\sum_{\ell} \|f_j\|_{L^{p_j}(w_j, 2^k Q_{\ell})}^{p_j} \right)^{p/p_j} \\
&\leq C 2^{-kn} \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}^p.
\end{aligned}$$

This completes our proof. □

For a family of balls $\{Q_k\}_k$ given by Proposition 2.6, for $s > 0$, we define the operator $\mathcal{M}_{\text{loc},s}$ by setting

$$\mathcal{M}_{\text{loc},s}(\vec{f}) = \sum_k \chi_{Q_k} \mathcal{M}_s(\vec{f} \chi_{\tilde{Q}_k}) \tag{6}$$

where $\tilde{Q}_j = 8Q_j$, $\vec{f} \chi_{\tilde{Q}_k} = (f_1 \chi_{\tilde{Q}_k}, \dots, f_m \chi_{\tilde{Q}_k})$, and the maximal function \mathcal{M}_s is defined by

$$\mathcal{M}_s(\vec{f})(x) = \sup_{Q \ni x} \prod_{j=1}^m \left(\frac{1}{|Q|} \int_Q |f_j(z)|^s dz \right)^{1/s}.$$

When $s = 1$, we drop the subindex s to write \mathcal{M}_{loc} instead of $\mathcal{M}_{\text{loc},1}$.

We have the following result.

Proposition 2.9 (i) If $p_1, \dots, p_m > s > 0$ and $\vec{w} \in A_{\vec{p}/s}^\infty$, then we have

$$\|\mathcal{M}_{\text{loc},s}(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

(ii) If $p_1, \dots, p_m \geq s > 0$ and $\vec{w} \in A_{\vec{p}/s}^\infty$, then we have

$$\|\mathcal{M}_{\text{loc},s}(\vec{f})\|_{L^{p,\infty}(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

Proof: The give the proof of (i) only. The proof of (ii) can be dealt by the analogous argument.

We first have

$$\begin{aligned} \int_{\mathbb{R}^n} |\mathcal{M}_{\text{loc},s}(\vec{f})(x)|^p \nu_{\vec{w}}(x) dx &= \sum_{\ell} \int_{Q_{\ell}} |\mathcal{M}_s(\vec{f} \chi_{\tilde{Q}_{\ell}})|^p \nu_{\vec{w}}(x) dx \\ &= \sum_{\ell} \int_{\mathbb{R}^n} |\mathcal{M}_s(\vec{f} \chi_{\tilde{Q}_{\ell}})|^p \nu_{\vec{w}_{\ell}}(x) dx \end{aligned}$$

where $\vec{w}_j = (w_1 \chi_{Q_j}, \dots, w_m \chi_{Q_j})$.

For each j , it can be verified that $\vec{w}_j \in A_{\vec{p}/s}^0$. Therefore, by Theorem 3.7 in [LOPTG], (ii) of Proposition 2.6 and Hölder inequality, we have

$$\begin{aligned} \int_{\mathbb{R}^n} |\mathcal{M}_{\text{loc},s}(\vec{f})(x)|^p \nu_{\vec{w}}(x) dx &\leq C \sum_{\ell} \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j, \tilde{Q}_{\ell})}^p \\ &\leq C \prod_{j=1}^m \left(\sum_{\ell} \|f_j\|_{L^{p_j}(w_j, \tilde{Q}_{\ell})}^{p_j} \right)^{p/p_j} \\ &\leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}^p. \end{aligned}$$

This completes our proof. □

When $m = 1$, we write $\widetilde{\mathcal{M}}_{\text{loc},s}$ instead of $\mathcal{M}_{\text{loc},s}$. As a direct consequence of Proposition 2.9, we have the following result.

Proposition 2.10 (i) If $p > s > 0$ and $w \in A_{p/s}^\infty$, then we have

$$\|\widetilde{\mathcal{M}}_{\text{loc},s}(f)\|_{L^p(w)} \leq C \|f\|_{L^p(w)}.$$

(ii) If $p = s$ and $w \in A_1^\infty$, then we have

$$\|\widetilde{\mathcal{M}}_{\text{loc},s}(\vec{f})\|_{L^{s,\infty}(w)} \leq C \|f\|_{L^s(w)}.$$

3 Main results

3.1 Weighted norm inequalities for multilinear operators

In this section, we establish the weighted norm inequalities for multilinear operators in the product of weighted L^p spaces by using the new class of weights $A_{\vec{p}}^\infty$.

Theorem 3.1 *Let T satisfy (H1), (H2) and (H3). Then the following statements hold:*

(i) *If $1 < p_1, \dots, p_m < \infty$ and $\vec{w} \in A_{\vec{p}}^\infty$, then*

$$\|T(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

(ii) *If $1 \leq p_1, \dots, p_m < \infty$ and at least one of the $p_j = 1$, then*

$$\|T(\vec{f})\|_{L^{p,\infty}(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

To prove Theorem 3.1, we need the following auxiliary propositions.

Proposition 3.2 *Let T satisfy (H1), (H2) and (H3). Then for any critical ball Q and $0 < \delta < 1/m$, we have*

$$\left(\frac{1}{|2Q|} \int_{2Q} |T(\vec{f})(z)|^\delta dz \right)^{1/\delta} \leq C \inf_{y \in 2Q} \mathfrak{M}(\vec{f})(y).$$

Proof: We split $f_j = f_j^0 + f_j^\infty$ where $f_j^0 = f_j \chi_{4Q}$ for all j . Then,

$$T(\vec{f})(x) = T(\vec{f}_0)(x) + \sum_{\alpha \in \mathcal{I}} T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x)$$

where $\vec{f}_0 = (f_1^0, \dots, f_m^0)$, $\alpha_1, \dots, \alpha_m \in \{0, \infty\}$ and $\mathcal{I} := \{\alpha = (\alpha_1, \dots, \alpha_m) : \alpha \neq (0, \dots, 0)\}$.

Therefore,

$$\begin{aligned} \left(\frac{1}{|2Q|} \int_{2Q} |T(\vec{f})(z)|^\delta dz \right)^{1/\delta} &\leq C \left(\frac{1}{|2Q|} \int_{2Q} |T(\vec{f}_0)(z)|^\delta dz \right)^{1/\delta} \\ &\quad + C \left(\frac{1}{|2Q|} \int_{2Q} \left| \sum_{\alpha \in \mathcal{I}} T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) \right|^\delta dz \right)^{1/\delta} \\ &:= I_1 + I_2. \end{aligned}$$

Note that

$$\|T(\vec{f}_0)\|_{L^{1/m,\infty}} \leq C \prod_{j=1}^m \|f_j^0\|_{L^1}.$$

This estimate together with Kolmogorov inequality implies

$$I_1 \leq \|T(\vec{f}_0)\|_{L^{1/m,\infty}(2Q, \frac{dz}{|2Q|})} \leq C \inf_{y \in 2Q} \mathfrak{M}(\vec{f})(y).$$

Let us take care I_2 . For each $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathcal{I}$ and $z \in 2Q$, we have, by (H1),

$$\begin{aligned}
|T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z)| &\leq C \int_{\mathbb{R}^{mn}} \frac{|f_1^{\alpha_1}(y_1) \dots f_m^{\alpha_m}(y_m)|}{(|z - y_1| + \dots + |z - y_m|)^{mn+N}} dy \\
&\leq C \sum_{k \geq 2} \int_{(2^{k+1}Q)^m \setminus (2^k Q)^m} \frac{|f_1(y_1) \dots f_m(y_m)|}{(|z - y_1| + \dots + |z - y_m|)^{mn+N}} dy \\
&\leq C \sum_{k \geq 2} 2^{-kN} \frac{1}{|2^{k+1}Q|^m} \int_{(2^{k+1}Q)^m} |f_1(y_1)| \dots |f_m(y_m)| dy \\
&\leq C \inf_{y \in 2Q} \mathfrak{M}(\vec{f})(y).
\end{aligned}$$

This completes our proof. □

Proposition 3.3 *Let T satisfy (H1), (H2) and (H3), let $0 < \delta < 1/m$. Then we have*

$$(M_{\text{loc},4}^\sharp |T(\vec{f})|^\delta(x))^{1/\delta} \leq C \mathfrak{M}(\vec{f})(x) + C \mathcal{M}_{\text{loc}}(\vec{f})(x).$$

Proof: It suffices to show that for each ball $B \ni x$ with $r_B \leq 4$, we can pick a number c_B so that

$$\left(\frac{1}{|B|} \int_B \left| |T(\vec{f})(z)|^\delta - |c_B|^\delta \right| dz \right)^{1/\delta} \leq C \mathfrak{M}(\vec{f})(x) + C \mathcal{M}_{\text{loc}}(\vec{f})(x).$$

By the same decomposition as in Proposition 3.2, we can write

$$\begin{aligned}
\left(\frac{1}{|B|} \int_B \left| |T(\vec{f})(z)|^\delta - |c_B|^\delta \right| dz \right)^{1/\delta} &\leq C \left(\frac{1}{|B|} \int_B |T(\vec{f})(z) - c_B|^\delta dz \right)^{1/\delta} \\
&\leq C \left(\frac{1}{|B|} \int_B |T(\vec{f}_0)(z)|^\delta dz \right)^{1/\delta} \\
&\quad + C \left(\frac{1}{|B|} \int_B \left| \sum_{\alpha \in \mathcal{I}} T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - c_B \right|^\delta dz \right)^{1/\delta} \\
&= E_1 + E_2.
\end{aligned}$$

By Kolmogorov inequality and the fact that T is bounded form $L^1 \times \dots \times L^1$ into $L^{1/m, \infty}$, we have

$$E_1 \leq C \prod_{j=1}^m \frac{1}{|4B|} \int_{4B} |f_j(z)| dz.$$

Let $\{Q_\ell\}_\ell$ be the family in Proposition 2.6. Then if $4B \cap Q_\ell \neq \emptyset$, we have $4B \subset \tilde{Q}_\ell := 8Q_\ell$. Therefore,

$$E_1 \leq C \mathcal{M}_{\text{loc}}(\vec{f})(x).$$

For the second term E_2 . Taking $c_B = \sum_{\alpha \in \mathcal{I}} T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x_B)$, we have

$$\sum_{\alpha \in \mathcal{I}} T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - c_B = \sum_{\alpha \in \mathcal{I}} \left[T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x_B) \right].$$

For each $\alpha \in \mathcal{I}$ and $N > 0$, by (H2), we write

$$\begin{aligned}
& \left| T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x_B) \right| \\
& \leq C \int_{\mathbb{R}^{mn}} |K(z, y_1, \dots, y_m) - K(x_B, y_1, \dots, y_m)| \prod_{j=1}^m |f_j^{\alpha_j}(y_j)| dy \\
& \leq C \sum_{k \geq 2} \int_{(2^{k+1}B)^m \setminus (2^k B)^m} |K(z, y_1, \dots, y_m) - K(x_B, y_1, \dots, y_m)| \prod_{j=1}^m |f_j(y_j)| dy \\
& \leq C \sum_{k \geq 2} \min\{1, (2^k r_B)^{-N}\} \frac{2^{-k\epsilon}}{|2^{k+1}B|^m} \int_{(2^{k+1}B)^m} \prod_{j=1}^m |f_j(y_j)| dy \\
& \leq C \sum_{k \geq k_0} \dots + C \sum_{k < k_0} \dots := E_{21} + E_{22}
\end{aligned}$$

where k_0 is the smallest integer so that $2^{k_0} r_B \geq 1$.

Let us estimate E_{22} first. Since $r_{2^{k+1}B} \leq 4$ for all $k < k_0$, the similar argument used in the estimate E_1 gives

$$\frac{1}{|2^{k+1}B|^m} \int_{(2^{k+1}B)^m} \prod_{j=1}^m |f_j(y_j)| dy \leq C \mathcal{M}_{\text{loc}}(\vec{f})(x)$$

for all $k < k_0$.

Therefore,

$$E_{22} \leq C \sum_{k < k_0} 2^{-k\epsilon} \mathcal{M}_{\text{loc}}(\vec{f})(x) \leq C \mathcal{M}_{\text{loc}}(\vec{f})(x).$$

For the term E_{21} , setting $\widehat{B} = 2^{k_0} B$, then $1 \leq r_{\widehat{B}} < 2$. We have

$$\begin{aligned}
E_{21} &= \sum_{k \geq k_0} (2^{k-k_0} 2^{k_0} r_B)^{-N} \frac{2^{-k\epsilon}}{|2^{k-k_0+1}\widehat{B}|^m} \int_{(2^{k-k_0+1}\widehat{B})^m} \prod_{j=1}^m |f_j(y_j)| dy \\
&= \sum_{k \geq k_0} 2^{-kN} \frac{1}{|2^k \widehat{B}|^m} \int_{(2^k \widehat{B})^m} \prod_{j=1}^m |f_j(y_j)| dy \\
&\leq C \mathfrak{M}(\vec{f})(x).
\end{aligned}$$

This completes our proof. □

We are now in position to prove Theorem 3.1.

Proof of Theorem 3.1: Since the proofs of (i) and (ii), we give only the proof of (i).

(i) Since $\nu_{\vec{w}} \in A_{mp}^\infty$, using Proposition 2.7 for $\delta < p < 1/m$, we have

$$\begin{aligned}
\|T(\vec{f})\|_{L^p(\nu_{\vec{w}})}^p &:= \|(T(\vec{f}))^\delta\|_{L^{p/\delta}(\nu_{\vec{w}})}^{p/\delta} \\
&\leq \int_{\mathbb{R}^n} |M_{\text{loc},\beta}(|T(\vec{f})|^\delta)(x)|^{p\delta} \nu_{\vec{w}}(x) dx \\
&\leq C \int_{\mathbb{R}^n} |M_{\text{loc},4}^\sharp(|T(\vec{f})|^\delta)(x)|^{p\delta} \nu_{\vec{w}}(x) dx + C \sum_{\ell} \nu_{\vec{w}}(2Q_\ell) \left(\frac{1}{2Q_k} \int_{2Q_k} |T(\vec{f})(z)|^\delta dz \right)^{p/\delta} \\
&= E_1 + E_2.
\end{aligned}$$

By Proposition 3.2, Proposition 2.6 and Proposition 2.8, we have

$$\begin{aligned}
E_2 &\leq C \sum_{\ell} \nu_{\vec{w}}(2Q_\ell) \left[\inf_{y \in 2Q_\ell} \mathfrak{M}(\vec{f})(y) \right]^p \\
&\leq C \sum_{\ell} \int_{Q_\ell} |\mathfrak{M}(\vec{f})(z)|^p \nu_{\vec{w}}(z) dz \\
&\leq C \|\mathfrak{M}(\vec{f})\|_{L^p(\nu_{\vec{w}})}^p \\
&\leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}^p.
\end{aligned}$$

It remains to estimate E_1 . In the light of Proposition 3.3, Proposition 2.8 and Proposition 2.9, we have

$$\begin{aligned}
E_1 &\leq C \int_{\mathbb{R}^n} |\mathfrak{M}(\vec{f})(z)|^p \nu_{\vec{w}}(z) dz + C \int_{\mathbb{R}^n} |\mathcal{M}_{\text{loc}}(\vec{f})(z)|^p \nu_{\vec{w}}(z) dz \\
&\leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}^p.
\end{aligned}$$

This completes our proof. □

3.2 Weighted norm inequalities for the commutators of multilinear operators with BMO_∞ functions

Let $\vec{b} = (b_1, \dots, b_m)$, where b_j is a locally integrable function for $j = 1, \dots, m$. We consider the m -linear commutator of T and \vec{b}

$$T_{\vec{b}}(\vec{f}) = \sum_{j=1}^m T_{\vec{b}}^j(\vec{f})$$

where

$$T_{\vec{b}}^j(\vec{f}) = b_j T(\vec{f}) - T(f_1, \dots, b_j f_j, \dots, f_m).$$

Let $\vec{\theta} = (\theta_1, \dots, \theta_m)$, $\theta_j \geq 0$ for all $j = 1, \dots, m$. For $\vec{b} \in BMO_{\vec{\theta}}$, we shall mean $b_j \in BMO_{\theta_j}$ for all $j = 1, \dots, m$ and $\|\vec{b}\|_{\vec{\theta}} = \sum_{j=1}^m \|b_j\|_{\theta_j}$.

Before coming to detail information, we need the following auxiliary results.

Proposition 3.4 *Let T satisfy (H1), (H2) and (H3) and $\vec{b} \in BMO_{\vec{\theta}}$. For any $p > 1$, the following holds for any critical ball Q and $0 < \delta < 1/m$,*

$$\left(\frac{1}{|2Q|} \int_{2Q} |T_{\vec{b}}(\vec{f})(z)|^{\delta} dz \right)^{1/\delta} \leq C \|\vec{b}\|_{\vec{\theta}} \inf_{y \in 2Q} \mathfrak{M}_p(\vec{f})(y).$$

Proof: By linearity it suffices to consider the commutator with one symbol as follows:

$$T_b(\vec{f}) = bT(\vec{f}) - T(bf_1, \dots, f_m), b \in BMO_{\theta}.$$

For any balls Q , we can write

$$T_b(\vec{f}) = (b - b_Q)T(\vec{f}) - T((b - b_Q)f_1, \dots, f_m) := I_1 + I_2.$$

Let $\delta < \delta' < 1/m$ and $1/s + 1/\delta' = 1/\delta$ so that $s > 1$. By Hölder inequality and Proposition 3.2, we have

$$\begin{aligned} I_1 &\leq \left(\frac{1}{|2Q|} \int_{2Q} |T(\vec{f})(z)|^{\delta'} dz \right)^{1/\delta'} \left(\frac{1}{|2Q|} \int_{2Q} |b(z) - b_Q|^s dz \right)^{1/s} \\ &\leq C \|b\|_{\theta} \inf_{y \in 2Q} \mathfrak{M}(\vec{f})(y) \\ &\leq C \|b\|_{\theta} \inf_{y \in 2Q} \mathfrak{M}_p(\vec{f})(y). \end{aligned}$$

To estimate I_2 , using the decomposition as in the proof of Proposition 3.2, we have

$$T((b - b_Q)f_1, \dots, f_m)(z) = T((b - b_Q)f_1^0, \dots, f_m^0)(z) + \sum_{\alpha \in \mathcal{I}} T((b - b_Q)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z)$$

Using Kolmogorov inequality and the fact that T maps continuously from $L^1 \times \dots \times L^1$ into $L^{1/m, \infty}$, we have

$$\begin{aligned} &\left(\frac{1}{|2Q|} \int_{2Q} |T((b - b_Q)f_1^0, \dots, f_m^0)(z)|^{\delta} dz \right)^{1/\delta} \\ &\leq \|T((b - b_Q)f_1^0, \dots, f_m^0)\|_{L^{1/m, \infty}(2Q, \frac{dz}{|2Q|})} \\ &\leq C \left(\frac{1}{|2Q|} \int_{4Q} |(b(z) - b_Q)f_1^0(z)| dz \right) \prod_{j \neq 1} \left(\frac{1}{|2Q|} \int_{4Q} |f_j^0(z)| dz \right) \\ &\leq C \left(\frac{1}{|2Q|} \int_{4Q} |(b(z) - b_Q)|^{p'} dz \right)^{1/p'} \prod_j \left(\frac{1}{|2Q|} \int_{4Q} |f_j^0(z)|^p dz \right)^{1/p} \\ &\leq C \|b\|_{\theta} \inf_{y \in 2Q} \mathfrak{M}_p(\vec{f})(y). \end{aligned}$$

For each $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathcal{I}$ and $z \in 2Q$, we have, by (H1), Hölder inequality and

Proposition 2.5,

$$\begin{aligned}
& |T((b - b_Q)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z)| \\
& \leq C \int_{\mathbb{R}^{mn}} \frac{|(b - b_Q)f_1^{\alpha_1}(y_1) \dots f_m^{\alpha_m}(y_m)|}{(|z - y_1| + \dots + |z - y_m|)^{mn+N}} dy \\
& \leq C \sum_{k \geq 2} \int_{(2^{k+1}Q)^m \setminus (2^kQ)^m} \frac{|(b - b_Q)f_1(y_1) \dots f_m(y_m)|}{(|z - y_1| + \dots + |z - y_m|)^{mn+N}} dy \\
& \leq C \sum_{k \geq 2} 2^{-kN} \left(\frac{1}{|2^{k+1}Q|} \int_{2^{k+1}Q} |(b - b_Q)f_1(y_1)| dy_1 \right) \prod_{j \neq 1} \left(\frac{1}{|2^{k+1}Q|} \int_{2^{k+1}Q} |f_j(y_j)| dy_j \right) \\
& \leq C \sum_{k \geq 2} 2^{-k(N-\theta)} \|b\|_\theta \left(\frac{1}{|2^{k+1}Q|} \int_{2^{k+1}Q} |f_1(y_1)|^p dy_1 \right)^{1/p} \prod_{j \neq 1} \left(\frac{1}{|2^{k+1}Q|} \int_{2^{k+1}Q} |f_j(y_j)| dy_j \right) \\
& \leq C \|b\|_\theta \inf_{y \in 2Q} \mathfrak{M}_p(\vec{f})(y).
\end{aligned}$$

This completes our proof. \square

Proposition 3.5 *Let T satisfy (H1), (H2) and (H3), let $0 < \delta < s < 1/m$ and $\vec{b} \in BMO_{\vec{\theta}}$. Then we have, for any $p > 1$,*

$$(M_{\text{loc},4}^\sharp(|T_{\vec{b}}(\vec{f})|^\delta)(x))^{1/\delta} \leq C \|\vec{b}\|_{\vec{\theta}} \left[\widetilde{M}_{\text{loc},s}(T(\vec{f}))(x) + \mathfrak{M}_p(\vec{f})(x) + \mathcal{M}_{\text{loc},p}(\vec{f})(x) \right]$$

for all $x \in \mathbb{R}^n$.

Proof: We need only to consider the commutator with one symbol as follows:

$$T_b(\vec{f}) = bT(\vec{f}) - T(bf_1, \dots, f_m)$$

where $b \in BMO_\theta$ and $\lambda \in \mathbb{R}$.

It suffices to show that for each ball $B \ni x$ with $r_B \leq 4$,

$$\left(\frac{1}{|B|} \int_B \left| |T_b(\vec{f})(z)|^\delta - |c_B|^\delta \right| dz \right)^{1/\delta} \leq C \mathfrak{M}(\vec{f})(x) + C \mathcal{M}_{\text{loc}}(\vec{f})(x)$$

where c_B is a constant which will be fixed later.

We write

$$T_b(\vec{f}) = (b - b_B)T(\vec{f}) - T((b - b_B)f_1, \dots, f_m)$$

By the same decomposition as in Proposition 3.2, we can write

$$\begin{aligned}
\left(\frac{1}{|B|} \int_B \left| |T_b(\vec{f})(z)|^\delta - |c_B|^\delta \right| dz \right)^{1/\delta} & \leq \left(\frac{1}{|B|} \int_B |T_b(\vec{f})(z) - c_B|^\delta dz \right)^{1/\delta} \\
& \leq C \left(\frac{1}{|B|} \int_B |(b - b_B)T(\vec{f})(z)|^\delta dz \right)^{1/\delta} \\
& \quad + C \left(\frac{1}{|B|} \int_B |T((b - b_B)f_1^0, \dots, f_m^0)(z)|^\delta dz \right)^{1/\delta} \\
& \quad + C \left(\frac{1}{|B|} \int_B \left| \sum_{\alpha \in \mathcal{I}} T((b - b_B)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - c_B \right|^\delta dz \right)^{1/\delta} \\
& = E_1 + E_2 + E_3.
\end{aligned}$$

By Hölder inequality and Proposition 2.5, we have

$$E_1 \leq C \|b\|_\theta \left(\frac{1}{|4B|} \int_B |T(\vec{f})(z)|^s dz \right)^{1/s}.$$

Repeating the argument as in Proposition 3.3, we conclude that

$$E_1 \leq C \|b\|_\theta \widetilde{\mathcal{M}}_{\text{loc},s}(T(\vec{f}))(x).$$

The arguments in Propositions 3.3 and 3.4 yields that

$$E_2 \leq C \|b\|_\theta \prod_{j=1}^m \left(\frac{1}{|4B|} \int_{4B} |f_j(z)|^p dz \right)^{1/p} \leq C \|b\|_\theta \mathcal{M}_{\text{loc},p}(\vec{f})(x).$$

For the second term E_3 . Taking $c_B = \sum_{\alpha \in \mathcal{I}} T((b - b_B)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x_B)$, we have

$$\begin{aligned} & \sum_{\alpha \in \mathcal{I}} T((b - b_B)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - c_B \\ &= \sum_{\alpha \in \mathcal{I}} \left[T((b - b_B)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - T(f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x_B) \right]. \end{aligned}$$

For each $\alpha \in \mathcal{I}$ and $N > 0$, by (H2), we write

$$\begin{aligned} & \left| T((b - b_B)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(z) - T((b - b_B)f_1^{\alpha_1}, \dots, f_m^{\alpha_m})(x_B) \right| \\ & \leq C \int_{\mathbb{R}^{mn}} |K(z, y_1, \dots, y_m) - K(x_B, y_1, \dots, y_m)| |(b - b_B)f_1^{\alpha_1}(y_1)| \prod_{j \neq 1}^m |f_j^{\alpha_j}(y_j)| dy \\ & \leq C \sum_{k \geq 2} \int_{(2^{k+1}B)^m \setminus (2^k B)^m} |K(z, y_1, \dots, y_m) - K(x_B, y_1, \dots, y_m)| \\ & \quad \times |(b - b_B)f_1(y_1)| \prod_{j \neq 1}^m |f_j(y_j)| dy \\ & \leq C \sum_{k \geq 2} \min\{1, (2^k r_B)^{-N}\} \frac{2^{-k\epsilon}}{|2^{k+1}B|^m} \int_{(2^{k+1}B)^m} |(b - b_B)f_1(y_1)| \prod_{j \neq 1}^m |f_j(y_j)| dy \\ & \leq C \sum_{k \geq k_0} \dots + C \sum_{k < k_0} \dots := E_{31} + E_{32} \end{aligned}$$

where k_0 is the smallest integer so that $2^{k_0} r_B \geq 1$.

Let us estimate E_{32} first. Since $r_{2^{k+1}B} \leq 4$ for all $k < k_0$, the similar argument used in the estimate E_2 gives

$$\frac{1}{|2^{k+1}B|^m} \int_{(2^{k+1}B)^m} |(b - b_B)f_1(y_1)| \prod_{j \neq 1}^m |f_j(y_j)| dy \leq C \|b\|_\theta \mathcal{M}_{\text{loc},p}(\vec{f})(x).$$

Therefore,

$$E_{32} \leq C \|b\|_\theta \sum_{k < k_0} 2^{-k\epsilon} \mathcal{M}_{\text{loc},p}(\vec{f})(x) \leq C \|b\|_\theta \mathcal{M}_{\text{loc},p}(\vec{f})(x).$$

For the term E_{31} , setting $\widehat{B} = 2^{k_0} B$, then $1 \leq r_{\widehat{B}} < 2$. We have

$$\begin{aligned}
E_{31} &= C \sum_{k \geq k_0} (2^{k-k_0} 2^{k_0} r_B)^{-N} \frac{2^{-k\epsilon}}{|2^{k-k_0+1} \widehat{B}|^m} \int_{(2^{k-k_0+1} \widehat{B})^m} |(b - b_B) f_1(y_1)| \prod_{j \neq 1}^m |f_j(y_j)| dy \\
&= C \sum_{k \geq k_0} 2^{-kN} \frac{1}{|2^k \widehat{B}|^m} \int_{(2^k \widehat{B})^m} |(b - b_B) f_1(y_1)| \prod_{j \neq 1}^m |f_j(y_j)| dy \\
&\leq C \sum_{k \geq k_0} 2^{-k(N-\theta)} \left(\frac{1}{|2^k \widehat{B}|^m} \int_{(2^k \widehat{B})^m} \prod_{j=1}^m |f_j(y_j)|^p dy \right)^{1/p} \\
&\leq C \|b\|_{\theta} \mathfrak{M}_p(\vec{f})(x).
\end{aligned}$$

This completes our proof. □

By similar arguments used in the proof of Theorem 3.1, we have the following result concerning the weighted norm inequality of the commutators $T_{\vec{b}}$.

Theorem 3.6 *Let T satisfy (H1), (H2) and (H3) and $\vec{b} \in BMO_{\vec{b}}$. Then we have, for $1 < p_1, \dots, p_m < \infty$ and $\vec{w} \in A_{\vec{p}}^\infty$,*

$$\|T_{\vec{b}}(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

4 Application to multilinear pseudodifferential operators

In this section, we will apply the obtained results to study the weighted norm inequalities for multilinear pseudodifferential operators.

Given a function $a : \mathbb{R}^{n(m+1)} \rightarrow \mathbb{C}$ satisfying certain growth conditions, we define the multilinear operator T_a to act on m functions $f_1, \dots, f_m \in \mathcal{S}(\mathbb{R}^n)$ by setting

$$T_a(\vec{f})(x) = \int_{\mathbb{R}^{mn}} a(x, \xi) \prod_{j=1}^m \hat{f}_j(\xi_j) e^{i\langle x, \xi_1 + \dots + \xi_m \rangle} d\xi$$

where $\xi = (\xi_1, \dots, \xi_m) \in \mathbb{R}^{mn}$.

We say that the symbol a belongs to the Hörmander class $m-S_{\rho, \delta}^l$, $l \in \mathbb{R}, \rho, \delta \in [0, 1]$, i.e., for all multi-indices $\alpha, \beta_1, \dots, \beta_m$ there holds

$$|\partial_x^\alpha \partial_{\xi_1}^{\beta_1} \dots \partial_{\xi_m}^{\beta_m} a(x, \xi)| \leq C(1 + |\xi_1| + \dots + |\xi_m|)^{m+\delta|\alpha|-\rho(|\beta_1|+\dots+|\beta_m|)}.$$

We have the following result.

Proposition 4.1 *Let $a \in m-S_{1, \delta}^0$, $0 \leq \delta < 1$. Then T_a satisfies (H1) and (H2).*

Proof: Let $\varphi_0 : \mathbb{R}^{mn} \rightarrow \mathbb{R}$ be a smooth radial function which is equal to 1 on the unit ball centered at origin and supported on its concentric double. Set $\varphi(\xi) = \varphi_0(\xi) - \varphi_0(2\xi)$ and $\varphi_k(\xi) = \varphi(2^{-k}\xi)$. Then, we have

$$\sum_{k=0}^{\infty} \varphi_k(\xi) = 1 \text{ for all } \xi \in \mathbb{R}^{mn}$$

and $\text{supp } \varphi_k \subset \{\xi : 2^{k-1} \leq |\xi| \leq 2^{k+1}\}$ for all $k \geq 1$. Moreover, for any multi-index α and $N \geq 0$, we have

$$|\partial_{\xi}^{\alpha} \varphi_k(\xi)| \leq c_{\alpha} 2^{-k|\alpha|}. \quad (7)$$

Then we can write

$$a(x, \xi) = \sum_{k=0}^{\infty} \varphi_k(\xi) a(x, \xi) := \sum_{k=0}^{\infty} a_k(x, \xi)$$

It is not difficult to show that T_a satisfies (H1). The proof of this part is standard and hence we omit details here.

It remains to check that T_a satisfies (H1). To do this, we will work with each component T_{a_k} , for $k \geq 0$. With the same notations as in condition (H2), we consider two cases:

Case 1: $j \neq 0$

In this situation, we can assume that $j = 1$ and $\max_k |y_k - y_1| := |y_0 - y_1|$.

Let $K_k(y_0, y_1, \dots, y_m)$ be the associated kernel of T_{a_k} . Then we have

$$K_k(y_0, y_1, \dots, y_m) = \int_{\mathbb{R}^{mn}} a_k(y_0, \xi) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi$$

Therefore,

$$\begin{aligned} & K_k(y_0, y_1, \dots, y_m) - K_k(y_0, y'_1, \dots, y_m) \\ &= \int_{\mathbb{R}^{mn}} a_k(y_0, \xi) \prod_{j \neq 1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} (e^{i\langle \xi_1, y_0 - y_1 \rangle} - e^{i\langle \xi_1, y_0 - y'_1 \rangle}) d\xi \end{aligned}$$

We consider two subcases:

Subcase 1.1: $|y'_1 - y_1| \geq 2^{-k}$

In this situation we have, for any integer $M \geq 0$,

$$\begin{aligned} & \left| K_k(y_0, y_1, \dots, y_m) - K_k(y_0, y'_1, \dots, y_m) \right| \\ & \leq \left| \int_{\mathbb{R}^{mn}} a_k(y_0, \xi) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ & \quad + \left| \int_{\mathbb{R}^{mn}} a_k(y_0, \xi) \prod_{j \neq 1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} e^{i\langle \xi_1, y_0 - y'_1 \rangle} d\xi \right| \\ & := E_1 + E_2. \end{aligned}$$

By integration by part, $|y_0 - y_1| \approx |y_0 - y'_1| \approx \sum_{k,l} |y_k - y_l|$, (7) and definition of the class $S_{1,0}^0$ we have

$$\begin{aligned} E_1 &\leq \sum_{|\alpha|=mn+N} |y_0 - y_1|^{-mn-M} \left| \int_{\mathbb{R}^{mn}} a_k(y_0, \xi) \partial_{\xi_1}^\alpha \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ &\leq \sum_{|\alpha|=mn+N} |y_0 - y_1|^{-mn-M} \left| \int_{\mathbb{R}^{mn}} \partial_{\xi_1}^\alpha a_k(y_0, \xi) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ &\leq C |y_0 - y_1|^{-mn-M} 2^{-k(mn+M-mn)}. \end{aligned}$$

Note that by interpolation the inequality above holds for all $M \geq 0$. Taking $M = N + \epsilon$, we have

$$\begin{aligned} E_1 &\leq C |y_0 - y_1|^{-mn-\epsilon-N} 2^{-k(N+\epsilon)} \\ &\leq C \frac{|y'_1 - y_1|^\epsilon}{(\sum_{k,l} |y_k - y_l|)^{mn+\epsilon}} |y_0 - y_1|^{-N}. \end{aligned}$$

Hence,

$$E_1 \leq C \frac{|y'_1 - y_1|^\epsilon}{(\sum_{k,l} |y_k - y_l|)^{mn+\epsilon}} \min\{1, |y_0 - y_1|^{-N}\} (2^k |y_1 - y'_1|)^{-\epsilon}.$$

Likewise,

$$E_2 \leq C \frac{|y'_1 - y_1|^\epsilon}{(\sum_{k,l} |y_k - y_l|)^{mn+\epsilon}} \min\{1, |y_0 - y_1|^{-N}\} (2^k |y_1 - y'_1|)^{-\epsilon}.$$

Hence, in this situation, T_a satisfies (H2).

Subcase 1.2: $|y'_1 - y_1| < 2^{-k}$

In this case, by integration by part, we write, for an integer $M \geq 0$,

$$\begin{aligned} &\left| K_k(y_0, y_1, \dots, y_m) - K_k(y_0, y'_1, \dots, y_m) \right| \\ &= \sum_{|\alpha|=M} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} a_k(y_0, \xi) (1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle}) \partial_{\xi_1}^\alpha \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ &= \sum_{|\alpha|=M} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} \partial_{\xi_1}^\alpha \left[a_k(y_0, \xi) (1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle}) \right] \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ &= \sum_{|\alpha|+|\beta|=M} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} \partial_{\xi_1}^\alpha a_k(y_0, \xi) \partial_{\xi_1}^\beta (1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle}) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right|. \end{aligned}$$

If $|\beta| = 0$, $|1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle}| \leq |\xi_1| |y_1 - y'_1| \leq 2^k |y_1 - y'_1|$. Hence

$$\begin{aligned} &\sum_{|\alpha|=M} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} \partial_{\xi_1}^\alpha a_k(y_0, \xi) (1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle}) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ &\leq C |y_0 - y_1|^{-M} 2^{-k(M-mn-1)} |y_1 - y'_1|. \end{aligned}$$

If $|\beta| > 0$, $|\partial_{\xi_1}^\beta (1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle})| \leq C|y_1 - y'_1|^{|\beta|}$. Hence

$$\begin{aligned} \sum_{|\alpha|+|\beta|=M, |\beta|>0} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} \partial_{\xi_1}^\alpha a_k(y_0, \xi) \partial_{\xi_1}^\beta (1 - e^{i\langle \xi_1, y_1 - y'_1 \rangle}) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ \leq C \sum_{|\alpha|=M} |y_0 - y_1|^{-M} 2^{-k(M-|\beta|-mn)} |y_1 - y'_1|^{|\beta|} \\ \leq C |y_0 - y_1|^{-M} 2^{-k(M-mn-1)} |y_1 - y'_1|. \end{aligned}$$

Hence, for any integer $M \geq 0$,

$$\left| K_k(y_0, y_1, \dots, y_m) - K_k(y_0, y'_1, \dots, y_m) \right| \leq C |y_0 - y_1|^{-M} 2^{-k(M-mn-1)} |y_1 - y'_1|.$$

By interpolation, this inequality holds for all $M \geq 0$. Taking $M = N + mn + \epsilon$, we have

$$\begin{aligned} \left| K_k(y_0, y_1, \dots, y_m) - K_k(y_0, y'_1, \dots, y_m) \right| \\ \leq C |y_0 - y_1|^{-N-mn-\epsilon} 2^{-k(N+\epsilon-1)} |y_1 - y'_1| \\ \leq C |y_0 - y_1|^{-N-mn-\epsilon} |y_1 - y'_1|^\epsilon (2^k |y_1 - y'_1|)^{1-\epsilon}. \end{aligned}$$

Therefore, in this situation, T_a satisfies (H2).

Case 2: $j = 0$

We can assume that $\max_k |y_k - y_1| := |y_0 - y_1|$. We also consider two subcases:

Subcase 2.1: $|y_0 - y'_0| > 2^{-k}$

By the similar argument to that of Subcase 1.1 we get that T_a satisfies (H2).

Subcase 2.2: $|y_0 - y'_0| \leq 2^{-k}$

In this case, we have

$$\begin{aligned} K_k(y_0, y_1, \dots, y_m) - K_k(y'_0, y_1, \dots, y_m) \\ = \int_{\mathbb{R}^{mn}} \left[a_k(y_0, \xi) \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} - a_k(y'_0, \xi) \prod_{j=1}^m e^{i\langle \xi_j, y'_0 - y_j \rangle} \right] d\xi \\ = \int_{\mathbb{R}^{mn}} \left[a_k(y_0, \xi) - a_k(y'_0, \xi) \right] \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \\ + \sum_{\ell=1}^m \int_{\mathbb{R}^{mn}} a_k(y'_0, \xi) \left[e^{i\langle \xi_\ell, y_0 - y_\ell \rangle} - e^{i\langle \xi_\ell, y'_0 - y_\ell \rangle} \right] \prod_{j<\ell} e^{i\langle \xi_j, y'_0 - y_j \rangle} \prod_{j>\ell} e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \\ = I + \sum_{\ell=1}^m I_\ell. \end{aligned}$$

Repeating the arguments as in Subcase 1.2, we get that for any $N \geq 0$

$$\sum_{\ell=1}^m |I_\ell| \leq C |y_0 - y_1|^{-N-mn-\epsilon} |y_0 - y'_0|^\epsilon (2^k |y_0 - y'_0|)^{1-\epsilon} \quad (8)$$

for some $\epsilon > 0$.

It remains to estimate I . We can write, for any integer $M \geq 0$,

$$\begin{aligned} |I| &\leq C \sum_{|\alpha|=M} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} [a_k(y_0, \xi) - a_k(y'_0, \xi)] \partial_{\xi_1}^\alpha e^{i\langle \xi_1, y_0 - y_1 \rangle} \prod_{j \neq 1} e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \\ &\leq C \sum_{|\alpha|=M} |y_0 - y_1|^{-M} \left| \int_{\mathbb{R}^{mn}} \partial_{\xi_1}^\alpha [a_k(y_0, \xi) - a_k(y'_0, \xi)] \prod_{j=1}^m e^{i\langle \xi_j, y_0 - y_j \rangle} d\xi \right| \end{aligned}$$

where in the last inequality we use integration by part.

By the Mean value Theorem and definition of the class $S_{1,\delta}^0$, we have

$$\left| \partial_{\xi_1}^\alpha [a_k(y_0, \xi) - a_k(y'_0, \xi)] \right| \leq C |y_0 - y'_0| 2^{-k|\alpha|+k\delta} = C |y_0 - y'_0| 2^{-kM+k\delta}.$$

Therefore, for any integer $M \geq 0$,

$$|I| \leq C |y_0 - y_1|^{-M} |y_0 - y'_0| 2^{-k(M-mn)+k\delta}. \quad (9)$$

By interpolation again, (9) still holds for any $M \geq 0$.

We now choose $\epsilon > 0$ so that $1 - \delta > \epsilon$. Taking $M = mn + N + \epsilon$, we have

$$|I| \leq C |y_0 - y_1|^{-(mn+N+\epsilon)} |y_0 - y'_0|^\epsilon 2^{-k(N-\delta-\epsilon+1)} (2^k |y - y_0|)^{1-\epsilon}. \quad (10)$$

The combination of (8) and (10) yields that T_a satisfies (H2). This completes our proof. \square

From Proposition 4.1, Theorem 3.1 and Theorem 3.6, we imply the following result.

Theorem 4.2 *Let $a \in m\text{-}S_{1,\delta}^0$, $0 \leq \delta < 1$. If T_a satisfies (H3), then the following statements hold:*

(i) *For $1 < p_1, \dots, p_m < \infty$ and $\vec{w} \in A_{\vec{p}}^\infty$, we have*

$$\|T_a(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)};$$

(ii) *If $1 \leq p_1, \dots, p_m < \infty$ and at least one of the $p_j = 1$, then*

$$\|T_a(\vec{f})\|_{L^{p,\infty}(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)};$$

(iii) *For any $\vec{b} \in BMO_{\vec{\theta}}$, $1 < p_1, \dots, p_m < \infty$ and $\vec{w} \in A_{\vec{p}}^\infty$, we have*

$$\|(T_a)_{\vec{b}}(\vec{f})\|_{L^p(\nu_{\vec{w}})} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(w_j)}.$$

We would like to give some relevant comments:

(i) It was proved in [GT] that if T_a be a multilinear pseudodifferential operator with the symbol a in the class $m-S_{1,\delta}^0$, $0 \leq \delta < 1$ and all of the transposes T_a^{*j} also have symbols in $m-S_{1,1}^0$, then T_a satisfies (H3). Hence, the conclusions in Theorem 4.2 hold for such a T_a .

(ii) In particular case when $m = 2$, the authors in [BT] proved that if T_a be a multilinear pseudodifferential operator with the symbol a in the class $2-S_{1,\delta}^0$, $\delta \in [0, 1)$ then all of the transposes T_a^{*1} and T_a^{*2} also have symbols in $2-S_{1,\delta}^0$, $\delta \in [0, 1)$. Therefore, by the previous remark, the results in Theorem 4.2 hold for T_a .

(iii) It can be believed that the obtained results in Theorem 4.2 still hold for multilinear pseudodifferential operators with the symbols $a \in m-S_{\rho,\delta}^l$ with $0 \leq \delta \leq \rho \leq 1$, $\delta < 1$, $0 < \rho$, $l < mn(\rho - 1)$. These results will be studied in the forth-coming paper.

References

- [BT] A. Bényi and R. Torres, Symbolic calculus and the transposes of bilinear pseudodifferential operators, *Comm. Partial Diff. Eq.* **28** (2003), 1161-1181.
- [B] T. A. Bui, Weighted norm inequalities for pseudo-differential operators and their commutators, preprint.
- [BHS1] B. Bongioanni, E. Haboure and O. Salinas, Commutators of Riezs transforms related to Schrödinger operators, *J. Fourier Anal. Appl.* **17** (2011), 115-134.
- [BHS2] B. Bongioanni, E. Haboure and O. Salinas, Classes of weights related to Schrödinger operators, *J. Math. Anal. Appl.* **373** (2011), 563-579.
- [CM1] R. Coifman and Y. Meyer, On commutators of singular integral and bilinear singular integrals, *Trans. Amer. Math. Soc.*, **212** (1975), 315-331.
- [CM2] R. Coifman and Y. Meyer, Au delà des opérateurs pseudodifférentiels, *Astérisque*, **57** (1978).
- [CM3] R. Coifman and Y. Meyer, Ondelettes et opérateurs, III, Hermann, Paris, 1990.
- [DZ] J. Dziubański and J. Zienkiewicz, Hardy spaces H^1 associated to Schrödinger operators with potential satisfying reverse Hölder inequality, *Rev. Mat. Iberoam.* **15** (1999), 279-296.
- [GT] L. Grafakos and R. Torres, Multilinear Calderón-Zygmund theory, *Adv. in Math.* **165** (2002), 124-164.
- [LOPTG] A.K. Lerner, S. Ombrosi, C. Pérez, R.T. Torres and R. Trujillo-González, New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory, *Adv. in Math.* **220** (2009), 1222-1264.
- [KS] C. Kenig and E. M. Stein, Multilinear estimates and fractional integration, *Math. Res. Lett.*, **6** (1999), 1-15.

- [T] L. Tang, Extrapolation from $A_{\infty}^{\rho, \infty}$, vector-valued inequalities and applications in the Schrödinger settings, preprint.